

Critically loaded multi-server queues with abandonments, retrials, and time-varying parameters

Young Myoung Ko and Natarajan Gautam

November 12, 2009

Abstract

In this paper, we consider modeling time-dependent multi-server queues that include abandonments and retrials. For the performance analysis of those, fluid and diffusion models called “strong approximations” have been widely used in the literature. Although they are proven to be asymptotically exact, their effectiveness as approximations in *critically loaded regimes* needs to be investigated. To that end, we find that existing fluid and diffusion approximations might be either inaccurate under simplifying assumptions or computationally intractable. To address that concern, this paper focuses on developing a methodology by adjusting the fluid and diffusion models so that they significantly improve the estimation accuracy. We illustrate the accuracy of our adjusted models by performing a number of numerical experiments.

1 Introduction

In this paper, we are interested in the precise analysis of time-varying many-server queues with abandonments and retrials described in Mandelbaum et al. [12] (See Figure 1). Inspired by call centers, there have been extensive studies on multi-server queues, especially having a large number of servers. Most of the recent studies utilize asymptotic analysis as it makes the problem tractable and also provides good approximations under certain conditions. Asymptotic analysis, typically, utilizes weak convergence to fluid and diffusion limits which is nicely summarized in Billingsley [2] and Whitt [18]. Methodologies to obtain fluid and diffusion limits, as described in Halfin and Whitt [6], have been developed in the literature using two different ways in terms of the traffic intensity.




Figure 1: Multi-server queue with abandonment and retrials, Mandelbaum et al. [12]

The first approach is to consider the convergence of a sequence of traffic intensities to a certain value. Relying on the value to which the sequence converges, there are three different operational regimes: efficiency driven (ED), quality and efficiency driven (QED), and quality driven (QD). Roughly speaking, if the traffic intensity (ρ) of the limit process is strictly greater than 1, it is called ED regime. If $\rho = 1$, then that is QED, otherwise QD. Many research studies have been done under the ED and QED regimes for multi-server queues like call centers (Halfin and Whitt [6], Puhalskii and Reiman [17], Garnet et al. [5], Whitt [19], Whitt [20], Pang and Whitt [16]). Recently, the QED regime, also known as “Halfin-Whitt regime”, has received a lot of attention; this is because it actually achieves both high utilization of servers and quality of service (Zeltyn and Mandelbaum [24]), and is a favorable operational regime for call centers with strict performance constraints (Mandelbaum and Zeltyn [14]).

The second way to obtain limit processes is to accelerate parameters keeping the traffic intensity fixed. An effective methodology called “uniform acceleration” or “strong approximations” which enables the analysis of time-dependent queues (Kurtz [9], Mandelbaum and Pats [10], Mandelbaum and Pats [11], Massey and Whitt [15], Whitt [21], Mandelbaum et al. [12], Hampshire et al. [7]) is included in this scheme and in fact is the basis of this paper.

The advantage of the strong approximations as described in Kurtz [9] is that it can be applied to a wide class of stochastic processes and can be nicely extended to time-dependent systems by combining with the results in Mandelbaum et al. [12]. However, it cannot be applied to multi-server queues directly due to an assumption that is not satisfied: i.e., for the diffusion model, the differentiability of the rate functions (e.g. net arrival rates and service rates) is necessary. But some rate functions are not differentiable everywhere since they are of the forms, $\min(\cdot, \cdot)$ or $\max(\cdot, \cdot)$. To extend the theory to non-smooth rate functions, Mandelbaum et al. [12] proves weak convergence by introducing a new derivative called “scalable Lipschitz derivative” and provides models for several queueing systems such as Jackson networks, multi-server queues with abandonments and retrials, multi-class preemptive priority queues, etc. In addition, several sets of differential equations are also provided to obtain the mean value and covariance matrix of the limit processes. It, however, turns out that the resulting sets of differential equations are computationally intractable to solve in general and hence the theorems cannot be applied to obtain numerical values of performance measures. In a follow-on paper, Mandelbaum et al. [13] provides numerical results for queue lengths and waiting times in multi-server queues with abandonments and retrials by adding an assumption to deal with computational intractability. Specifically, the paper assumes measure zero at a set of time points where the fluid model hits non-differentiable points, which eventually enables us to apply Kurtz’s diffusion models. However, as pointed out in Mandelbaum et al. [13], if the system stays close to a critically loaded phase for a long time (i.e. *lingering* around a non-differentiable point), their approach may cause significant inaccuracy.

To explain this inaccuracy in detail, consider a multi-server queue with abandonments and retrials as shown in Figure 1. As an example we select numerical values $n_t = 50, \mu_t^1 = 1, \mu_t^2 = 0.2$ for all t , whereas λ_t alternates between $\lambda_t^1 = 45$ and $\lambda_t^2 = 55$ every two units of time (the parameters are defined in Section 2 and illustrated in Figure 1). Using the measure-zero assumption in Mandelbaum et al. [13], we graph $E[x_1(t)]$ and $E[x_2(t)]$ in Figure 2 (a), and also $Var[x_1(t)]$, $Var[x_2(t)]$, and $Cov[x_1(t), x_2(t)]$ in Figure 2 (b). Notice that, although $E[x_1(t)]$ is reasonably accurate, the others ($E[x_2(t)]$, $Var[x_1(t)]$, $Var[x_2(t)]$, and $Cov[x_1(t), x_2(t)]$) are not accurate at all. The reason for that is the system lingers around the non-differentiable points. In addition, if one were to solve the differential equations numerically using computationally intractable techniques via the Lipschitz derivatives as described in Mandelbaum et al. [12], the similar level of inaccuracy occurs. We explain

this in detail in Section 3.2. However, this does nourish the need for a methodology to accurately predict the system performance which is the focus of this study.

Having motivated the need to develop a methodology for the critically loaded phase, we now describe its importance. According to Mandelbaum and Pats [11] and Mandelbaum et al. [13], time-dependent queues make transitions among three phases: underloaded, critically loaded, and overloaded. The phase of the system is determined by the fluid model. The limit process in the strong approximations does not require any regimes such as QD, QED, or ED. However, from Section 1.4 in Zeltyn and Mandelbaum [24], we could find a rough correspondence between the operational regimes (QD, QED, and ED) and the phases in time-varying queues (underloaded, critically loaded, and overloaded). Recall that the QED regime is favorable to the operation of the call centers. Therefore, capturing the dynamics of multi-server queues in the critically loaded phase is also of significant importance. Nonetheless, from Figure 2, we found two major issues in the existing approach: 1) the fluid model (where the non-differentiability issue is actually irrelevant) is itself inaccurate and 2) sharp spikes which cause massive estimation errors are observed at the non-differentiable points in the diffusion model in contrast to the smooth curves in the simulation. In this paper, we approach the above two issues from a different point of view and provide an effective solution to them. Considering those, the contributions of this paper can be summarized as follows:

1. To the best of our knowledge, inaccuracy in the fluid model has never been addressed in the literature. We explain why it happens and ameliorate the fluid model.
2. Sharp spikes observed in the diffusion model cannot be resolved using the methodology in the literature. We provide a reasonable approximation-methodology so that it could smoothen the spikes and improve the estimation accuracy dramatically.

We now describe the organization of this paper. In Section 2, we state the problem considered in this paper. In Section 3, we summarize the strong approximations in Kurtz [9] and Mandelbaum et al. [12], and describe the above issues in detail. In Section 4, we construct an adjusted fluid model to estimate the exact mean value of the system state. However, this would not immediately result in a computationally feasible approach. For that, in Section 5, we explain our Gaussian-based approximations to achieve computational feasibility and smoothness in the diffusion model. Further





Figure 2: Simulation vs Fluid and diffusion model with measure-zero assumption

investigation on the adjusted models is provided in Section 6 to show how actually our adjusted models contribute to the estimation accuracy. In Section 7, we provide a number of numerical examples and compare against the existing approach as well as simulation. Finally, in Section 8, we make concluding remarks and explain directions for future work.

2 Problem description

Consider Figure 1 that illustrates a multi-server queue with abandonments and retrials as described in Mandelbaum et al. [12] and Mandelbaum et al. [13]. There are n_t number of servers in the service node at time t . Customers arrive to the service node according to a non-homogeneous Poisson process at rate λ_t . The service time of each customer follows a distribution having a memoryless property at rate μ_t^1 . Customers in the queue are served under the FCFS policy and the abandonment rate of customers is β_t with exponentially distributed time to abandon. Abandoning customers leave the system with probability p_t or go to a retrial queue with probability $1 - p_t$. The retrial queue is equivalent to an infinite-server-queue and hence each customer in the retrial queue waits there for a random amount of time with mean $1/\mu_t^2$ and returns to the service node.

Let $X(t) = (x_1(t), x_2(t))$ be the system state where $x_1(t)$ is the number of customers in the service node and $x_2(t)$ is the number of customers in the retrial queue. Then, $X(t)$ is the unique solution to the following integral equations:

$$\begin{aligned} x_1(t) = & x_1(0) + Y_1\left(\int_0^t \lambda_s ds\right) + Y_2\left(\int_0^t x_2(s) \mu_s^2 ds\right) - Y_3\left(\int_0^t (x_1(s) \wedge n_s) \mu_s^1 ds\right) \\ & - Y_4\left(\int_0^t (x_1(s) - n_s)^+ \beta_s (1 - p_s) ds\right) - Y_5\left(\int_0^t (x_1(s) - n_s)^+ \beta_s p_s ds\right), \end{aligned} \quad (1)$$

$$x_2(t) = x_2(0) + Y_4\left(\int_0^t (x_1(s) - n_s)^+ \beta_s (1 - p_s) ds\right) - Y_2\left(\int_0^t x_2(s) \mu_s^2 ds\right), \quad (2)$$

where Y_i 's are independent rate-1 Poisson processes.

The performance measures we are interested in are $E[X(t)]$ and $Cov[X(t), X(t)]$ (i.e. $Var[x_1(t)]$, $Var[x_2(t)]$, and $Cov[x_1(t), x_2(t)]$) for any given time $t \in [0, T]$, where $T < \infty$ is a constant. Especially, we have an interest in the system that is *lingering* near the critically loaded phase for a long time. Anyhow, as one may notice, the above two equations (1) and (2) cannot be solved directly. If all the parameters are constant, i.e. $\lambda_t = \lambda$, $\mu_t^1 = \mu^1$, $\mu_t^2 = \mu^2$, $\beta_t = \beta$, and $p_t = p$, one can consider

a Continuous Time Markov Chain (CTMC) model to obtain the performance measures. However, even assuming constant parameters, calculating the performance measures at any given time t is hard since $x_1(t)$ and $x_2(t)$ both are unbounded and solving balance equations in the two or more dimensional spaces requires tremendous efforts. Furthermore, when the number of servers is large, computational issues might arise. We, accordingly, would try to take advantage of an asymptotic methodology that is adequate for the analysis of time-varying systems with large number of servers. Nevertheless, as briefly mentioned in Section 1, we found that the existing methodologies are either computationally intractable or significantly inaccurate in the critically loaded phase. The objective of this paper is to develop a new approach to enhance the accuracy in estimating the mean value and covariance matrix for the multi-server queues with abandonments and retrials.

To do so, we start by summarizing the strong approximations and addressing the potential limitations in the following section.

3 Summary of the strong approximations

In Section 3.1, we recapitulate the strong approximations in Kurtz [9] and Mandelbaum et al. [12]. In Section 3.2, we explain what produces estimation errors and why existing methodologies do not fix them.

3.1 Strong approximations

In this section, we review the fluid and diffusion approximations developed by Kurtz [9] that we would leverage upon for our methodology. We also briefly mention the result in Mandelbaum et al. [12] which extends Kurtz's result to models involving non-smooth rate functions. Moreover, it is worthwhile to note that for $n \in \mathbf{N}$, the state of the queueing system $X_n(t)$ includes jumps but the limit process is continuous. Therefore, the weak convergence result that is presented is with respect to uniform topology in Space D (Billingsley [2] and Whitt [18]).

Let $X(t)$ be an arbitrary d -dimensional stochastic process which is the solution to the following integral equation:

$$X(t) = x_0 + \sum_{i=1}^k l_i Y_i \left(\int_0^t f_i(s, X(s)) ds \right), \quad (3)$$

where $x_0 = X(0)$ is a constant, Y_i 's are independent rate-1 Poisson processes, $l_i \in \mathbf{Z}^d$ for $i \in \{1, 2, \dots, k\}$ are constant, and f_i 's are continuous functions such that $|f_i(t, x)| \leq C_i(1 + |x|)$ for some $C_i < \infty$, $t \leq T$ and $T < \infty$. Note that we just consider a finite number of l_i 's to simplify proofs, which is reasonable for real world applications.

Notice that a special case of $X(t)$ in equation (3) is the $X(t)$ we described in equations (1) and (2) in our problem explained in Section 2. Following the notation in equation (3), we have, for our problem in Section 2, $x = (x_1, x_2)$ and $t \leq T$,

$$\begin{aligned} f_1(t, x) &= \lambda_t, f_2(t, x) = \mu_t^2 x_2, f_3(t, x) = \mu_t^1 (x_1 \wedge n_t), \\ f_4(t, x) &= \beta_t(1 - p_t)(x_1 - n_t)^+, f_5(t, x) = \beta_t p_t (x_1 - n_t)^+, \\ l_1 &= \begin{pmatrix} 1 \\ 0 \end{pmatrix}, l_2 = \begin{pmatrix} 1 \\ -1 \end{pmatrix}, l_3 = \begin{pmatrix} -1 \\ 0 \end{pmatrix}, l_4 = \begin{pmatrix} -1 \\ 1 \end{pmatrix}, \text{ and } l_5 = \begin{pmatrix} -1 \\ 0 \end{pmatrix}. \end{aligned}$$

Coming back to the generalized $X(t)$ process, we reiterate that it is usually not tractable to solve the integral equation (3). Therefore, to approximate the $X(t)$ process, define a sequence of stochastic processes $\{X_n(t)\}$ which satisfy the following integral equation:

$$X_n(t) = x_0 + \sum_{i=1}^k \frac{1}{n} l_i Y_i \left(\int_0^t n f_i(s, X_n(s)) ds \right).$$

Typically the process $X_n(t)$ (usually called a scaled process) is obtained by taking n times faster rates of events and $1/n$ of the increment of the system state. This type of setting is used in the literature and is denoted as “uniform acceleration” in Massey and Whitt [15], Mandelbaum et al. [12], and Mandelbaum et al. [13]. Then, the following theorem provides the fluid model to which $\{X_n(t)\}$ converges almost surely as $n \rightarrow \infty$. Define

$$F(t, x) = \sum_{i=1}^k l_i f_i(t, x). \tag{4}$$

Theorem 1 (Fluid model, Kurtz [9]). *If there is a constant $M < \infty$ such that $|F(t, x) - F(t, y)| \leq M|x - y|$ for all $t \leq T$ and $T < \infty$. Then, $\lim_{n \rightarrow \infty} X_n(t) = \bar{X}(t)$ a.s. where $\bar{X}(t)$ is the solution to*

the following integral equation:

$$\bar{X}(t) = x_0 + \sum_{i=1}^k l_i \int_0^t f_i(s, \bar{X}(s)) ds.$$

Note that $\bar{X}(t)$ is a deterministic time-varying quantity. We will subsequently connect $\bar{X}(t)$ and $X(t)$ defined in equation (3), but before that we provide the following result. Once we have the fluid model, we can obtain the diffusion model from the scaled centered process $(D_n(t))$. Define $D_n(t)$ to be $\sqrt{n}(X_n(t) - \bar{X}(t))$. Then, the limit process of $D_n(t)$ is provided by the following theorem.

Theorem 2 (Diffusion model, Kurtz [9]). *If f_i 's and F , for some $M < \infty$, satisfy*

$$|f_i(t, x) - f_i(t, y)| \leq M|x - y| \quad \text{and} \quad \left| \frac{\partial}{\partial x_i} F(t, x) \right| \leq M, \quad \text{for } i \in \{1, \dots, k\} \text{ and } 0 \leq t \leq T,$$

then $\lim_{n \rightarrow \infty} D_n(t) = D(t)$ where $D(t)$ is the solution to

$$D(t) = \sum_{i=1}^k l_i \int_0^t \sqrt{f_i(s, \bar{X}(s))} dW_i(s) + \int_0^t \partial F(s, \bar{X}(s)) D(s) ds,$$

$W_i(\cdot)$'s are independent standard Brownian motions, and $\partial F(t, x)$ is the gradient matrix of $F(t, x)$ with respect to x .

Remark 1. *Theorem 2 requires that $F(\cdot, \cdot)$ has a continuous gradient matrix. Therefore, if we don't have such an F , then we cannot apply Theorem 2 directly to obtain the diffusion model.*

Remark 2. *According to Ethier and Kurtz [3], if $D(0)$ is a constant or a Gaussian random vector, then $D(t)$ is a Gaussian process.*

Now, we have the fluid and diffusion models for $X_n(t)$. Therefore, for a large n , $X_n(t)$ is approximated by

$$X_n(t) \approx \bar{X}(t) + \frac{D(t)}{\sqrt{n}}.$$

If we follow this approximation, we can also approximate the mean and covariance matrix of $X_n(t)$

denoted by $E[X_n(t)]$ and $Cov[X_n(t), X_n(t)]$ respectively as

$$E[X_n(t)] \approx \bar{X}(t) + \frac{E[D(t)]}{\sqrt{n}}, \quad (5)$$

$$Cov[X_n(t), X_n(t)] \approx \frac{Cov[D(t), D(t)]}{n}. \quad (6)$$

In equations (5) and (6), only $\bar{X}(t)$ is known. Therefore, in order to get approximated values of $E[X_n(t)]$ and $Cov[X_n(t), X_n(t)]$, we need to obtain $E[D(t)]$ and $Cov[D(t), D(t)]$. The following theorem provides a methodology to obtain $E[D(t)]$ and $Cov[D(t), D(t)]$.

Theorem 3 (Mean and covariance matrix of linear stochastic systems, Arnold [1]). *Let $Y(t)$ be the solution to the following linear stochastic differential equation.*

$$dY(t) = A(t)Y(t)dt + B(t)dW(t), \quad Y(0) = 0,$$

where $A(t)$ is a $d \times d$ matrix, $B(t)$ is a $d \times k$ matrix, and $W(t)$ is a k -dimensional standard Brownian motion. Let $M(t) = E[Y(t)]$ and $\Sigma(t) = Cov[Y(t), Y(t)]$. Then, $M(t)$ and $\Sigma(t)$ are the solution to the following ordinary differential equations:

$$\begin{aligned} \frac{d}{dt}M(t) &= A(t)M(t) \\ \frac{d}{dt}\Sigma(t) &= A(t)\Sigma(t) + \Sigma(t)A(t)' + B(t)B(t)'. \end{aligned} \quad (7)$$

Corollary 1. *If $M(0) = 0$, then $E[M(t)] = 0$ for $t \geq 0$.*

By Corollary 1, if $D(0) = 0$, then $E[D(t)] = 0$ for $t \geq 0$. Therefore, if $\bar{X}(0) = X(0) = x_0$, then we can rewrite (5) to be

$$E[X_n(t)] \approx \bar{X}(t).$$

Recalling Remark 1, the diffusion model in Kurtz [9] requires differentiability of rate functions. Otherwise, we cannot apply Theorem 2. To get this problem under control, Mandelbaum et al. [12] introduces a new derivative called “scalable Lipschitz derivative” and proves weak convergence using it. Unlike the result in Kurtz [9], it turns out that the diffusion limit may not be a Gaussian

process when rate functions are not differentiable everywhere. In Mandelbaum et al. [12], expected values of the diffusion model may not be zero (compare it with Corollary 1) and could adjust the inaccuracy in the fluid model (see Mandelbaum et al. [13]). The resulting differential equations for the diffusion model, however, are computationally intractable. For example, in Mandelbaum et al. [12], one of the differential equations has the following form:

$$\begin{aligned} \frac{d}{dt}E[Q_1^{(1)}(t)] &= (\mu_t^1 \mathbf{1}_{\{Q_1^{(0)} \leq n_t\}} + \beta_t \mathbf{1}_{\{Q_1^{(0)} > n_t\}})E[Q_1^{(1)}(t)^-] \\ &\quad - (\mu_t^1 \mathbf{1}_{\{Q_1^{(0)} < n_t\}} + \beta_t \mathbf{1}_{\{Q_1^{(0)} \geq n_t\}})E[Q_1^{(1)}(t)^+] + \mu_t^2 E[Q_2^{(1)}(t)], \end{aligned} \quad (8)$$

rendering it to be intractable.

Therefore, Mandelbaum et al. [13], as we understand, resorts to the method in Kurtz [9] by assuming measure zero at non-smooth points to avoid computational difficulty.

3.2 Inaccuracy of strong approximations

Though not mentioned in any previous studies, to the best of our knowledge, the fluid model has the possibility of being inaccurate when approximating the mean value of the system state. Consider the actual integral equation to get the exact value of $E[X(t)]$ by the following theorem.

Theorem 4 (Expected value of $X(t)$). *Consider $X(t)$ defined in equation (3). Then, for $t \leq T$, $E[X(t)]$ is the solution to the following integral equation.*

$$E[X(t)] = x_0 + \sum_{i=1}^k l_i \int_0^t E[f_i(s, X(s))] ds \quad (9)$$

Proof. Take expectation on both sides of equation (3). Then,

$$\begin{aligned} E[X(t)] &= x_0 + \sum_{i=1}^k l_i E \left[Y_i \left(\int_0^t f_i(s, X(s)) ds \right) \right] \\ &= x_0 + \sum_{i=1}^k l_i E \left[\int_0^t f_i(s, X(s)) ds \right] \text{ since } Y_i(\cdot)\text{'s are non-homogeneous Poisson processes} \\ &= x_0 + \sum_{i=1}^k l_i \int_0^t E[f_i(s, X(s))] ds \text{ by Fubini theorem in Folland [4].} \end{aligned}$$

Therefore, we prove the theorem. $\square$

Comparing Theorems 1 and 4, notice that we cannot conclude that $\bar{X}(t)$ in Theorem 1 and $E[X(t)]$ in Theorem 4 are close enough since $E[f_i(t, X(t))] \neq f_i(t, E[X(t)])$. In some applications, f_i 's might be constants or linear combinations of components of $X(t)$. In those cases, Theorem 4 and the following corollary imply that the fluid model would be the exact mean value of the system state.

Corollary 2. *If $f_i(t, x)$'s are constants or linear combinations of the components of x , Then,*

$$E[X(t)] = \bar{X}(t),$$

where $X(t)$ is the solution to (3) and $\bar{X}(t)$ is the deterministic fluid model from theorem 1.

Proof. Using linearity of expectation in Williams [23], we can obtain the same integral equation for both $E[X(t)]$ and $\bar{X}(t)$. $\square$

However, if we have different forms of f_i 's where $E[f_i(t, X(t))] \neq f_i(t, E[X(t)])$, then the fluid model would be inaccurate. Notice that the fluid model does not require the differentiability of rate functions in both Kurtz [9] and Mandelbaum et al. [12]. Therefore, in this problem, the differentiability issue in rate functions is actually irrelevant.

Now we move our attention to inaccuracy in the diffusion model. We use the annotated version of Figure 2 (via Figure 3) here for the clear explanation. Figures 3 (a) and (b) show the mean value and covariance matrix of the system against those of the simulation respectively. Since the number of servers is 50, as shown in Figure 3 (a), the mean value of $x_1(t)$ is fluctuating near the critically loaded point. From the figure, we also confirm that the fluid model is quite inaccurate for the mean value of $x_2(t)$. For the covariance matrix, as shown in Figure 3 (b), the diffusion model brings about immense estimation errors (sharp spikes) in the vicinity of the critically loaded time points. Notice that from Figure 3 (b) we found that *even if the differential equations such as equation (8) in Mandelbaum et al. [12], which are known to be true, can be numerically solvable, it does not contribute to improving the estimation accuracy*. In the figure, the time point t_0 is the time when the fluid model hits a critically loaded point for the first time. *The differential equations in Mandelbaum et al. [12] are virtually same as those in Mandelbaum et al. [13] which assume measure*





Figure 3: Simulation vs Fluid and diffusion model with measure-zero assumption




Figure 4: Construction of a new process

zero for the computational tractability until the fluid model reaches a critically loaded point for the first time. Therefore, we can think that the graphs before time t_0 in Figure 3 are exactly same as those obtained from the methodology in Mandelbaum et al. [12] though we could not get the graphs after t_0 . However, as seen in Figure 3 (b), the estimation errors become apparent much earlier than the time point t_0 . Therefore, we figure out that the methodology in Mandelbaum et al. [12] does not remove the sharp spikes at least until the time t_0 . Moreover, from the shapes of the differential equations, we would conjecture that the methodology in Mandelbaum et al. [12] might not get rid of the sharp spikes even after the time t_0 . The drift matrix of the diffusion model in Mandelbaum et al. [12] still makes sudden changes at the critically loaded point which actually causes the spikes. We will revisit and explain it in Section 6.

In the next two sections, we describe our approach to the above issues in both fluid and diffusion models. In Section 4, we address the inaccuracy in the fluid model by a constructing new process. In particular, in Section 5, based on the adjusted fluid model, we explain how to remove the sharp spikes that causes vast estimation errors in the diffusion model.

4 Adjusted fluid model

The basic idea of our approach is to construct a new process, $Z(t)$, so that its fluid model is exactly the same as the mean value of the original process $X(t)$ as described in Theorem 4 (this is schematically explained in Figure 4). Although we concentrate on multi-server queues, this approach can be applied to more general types of stochastic systems. Therefore, we borrow the more general notation in Section 3.1 (as opposed to that in Section 2).

To begin with, define a set $\mathbb{F}$ of all distribution functions that have a finite mean and covariance matrix in $\mathbf{R}^d$. This set is valid for the fluid model since conditions on f_i 's guarantee that $E[|X(t)|] < \infty$ and $|Cov[X(t), X(t)]| < \infty$ for all $t \leq T$. Define a subset $\mathbb{F}_0$ of $\mathbb{F}$ such that any $h \in \mathbb{F}_0$ has zero mean. We call an element of $\mathbb{F}_0$ a “base distribution” for the remainder of this paper.

Proposition 1. *For $t \leq T$ and $i \in 1, 2, \dots, k$, let $\mu(t) = E[X(t)]$. Then, $E[f_i(t, X(t))]$ can be represented as a function of $\mu(t)$, i.e., there exists a function $g_i(t, \cdot)$ such that*

$$g(t, \mu(t)) = E[f_i(t, X(t))].$$

Proof. For fixed $t_0 \leq T$, suppose the distribution of $X(t_0)$ is F . Then, $F \in \mathbb{F}$. For $F \in \mathbb{F}$, we can always find $F_0 \in \mathbb{F}_0$ such that $F(x) = F_0(x - \mu)$ where $\mu = E[X(t_0)] = \int_{\mathbf{R}^d} x dF$. Then,

$$\begin{aligned} E[f_i(t_0, X(t_0))] &= \int_{\mathbf{R}^d} f_i(t_0, x) dF \\ &= \int_{\mathbf{R}^d} f_i(t_0, x + \mu) dF_0. \end{aligned}$$

Since the integration removes x , by making t_0 and μ variables (i.e. substitute t_0 and μ with t and $\mu(t)$ respectively), we have

$$E[f_i(t, X(t))] = g_i(t, \mu(t)), \text{ for some function } g_i.$$

□

Remark 3. *Proposition 1 does not mean that $\mu(\cdot)$ completely identifies the function $g_i(\cdot, \cdot)$. In fact, the function $g_i(\cdot, \cdot)$ might be unknown unless the base distribution is identified but we can say that such a function $g_i(\cdot, \cdot)$ exists.*

For $t \leq T$, let $\mu(t) = E[X(t)]$. Let $g_i(t, \mu(t)) = E[f_i(t, X(t))]$ for $i \in \{1, \dots, k\}$. Then, we can construct a new stochastic process $Z(t)$ which is the solution to the following integral equation:

$$Z(t) = z_0 + \sum_{i=1}^k l_i Y_i \left(\int_0^t g_i(s, Z(s)) ds \right). \quad (10)$$

Based on equation (10), define a sequence of stochastic processes $\{Z_n(t)\}$ satisfying

$$Z_n(t) = x_0 + \sum_{i=1}^k \frac{1}{n} l_i Y_i \left(\int_0^t n g_i(s, Z_n(s)) ds \right). \quad (11)$$

Next, we would like to obtain the fluid model for $Z_n(t)$. Before doing that, we need to check whether the functions g_i 's satisfy the conditions to apply Theorem 1. Following lemmas show that g_i 's actually meet those conditions. The proofs of the lemmas are provided in Appendix A.

Lemma 1. *If $|f_i(t, x)| \leq C_i(1 + |x|)$ for $t \leq T$, then $g_i(t, x)$'s satisfy*

$$|g_i(t, x)| \leq D_i(1 + |x|) \quad \text{for some } D_i < \infty.$$

For the next lemma, we would like to define

$$G(t, x) = \sum_{i=1}^k l_i g_i(t, x). \quad (12)$$

Lemma 2. *For $t \leq T$, if $|f_i(t, x) - f_i(t, y)| \leq M|x - y|$, then $g_i(t, x)$'s satisfy*

$$|g_i(t, x) - g_i(t, y)| \leq M|x - y|,$$

and if $|F(t, x) - F(t, y)| \leq M|x - y|$, then $G(t, x)$ satisfies

$$|G(t, x) - G(t, y)| \leq M|x - y|.$$

Lemmas 1 and 2 show that if f_i 's satisfy the conditions to obtain the fluid limit of $X_n(t)$, then g_i 's are also eligible for the fluid model of $Z_n(t)$. Therefore, we are now able to provide the adjusted fluid model based on Lemmas 1 and 2.

Theorem 5 (Adjusted fluid model). *Assume*

$$|f_i(t, x)| \leq C_i(1 + |x|) \quad \text{for } i \in \{1, \dots, k\}, \quad (13)$$

$$|F(t, x) - F(t, y)| \leq M|x - y|. \quad (14)$$

Then, $\lim_{n \rightarrow \infty} Z_n(t) = \bar{Z}(t)$ a.s., where $\bar{Z}(t)$ is the solution to the following integral equation:

$$\bar{Z}(t) = x_0 + \sum_{i=1}^k l_i \int_0^t g_i(s, \bar{Z}(s)) ds, \quad (15)$$

and furthermore

$$\bar{Z}(t) = E[X(t)] = x_0 + \sum_{i=1}^k l_i \int_0^t E[f_i(s, X(s))] ds. \quad (16)$$

Proof. From Lemmas 1 and 2, (13) and (14) imply

$$|g_i(t, x)| \leq D_i(1 + |x|) \quad \text{and} \quad |G(t, x) - G(t, y)| \leq M|x - y|.$$

Therefore, by Theorem 1, we have equation (15), and by the definition of $g_i(t, x)$'s, we have equation (16). $\square$

Comparing equation (16) with equation (9) in Theorem 4, we notice that Theorem 5 via equation (16) could provide the exact estimation of $E[X(t)]$. Though Theorem 5 provides the exact estimation of $E[X(t)]$, the functions g_i 's cannot be identified unless the base distribution is known, which forces us to develop an algorithm to find g_i 's. Nonetheless, when applying our adjusted fluid model to the multi-server queues with abandonments and retrials, we, in fact, have a good candidate distribution to obtain g_i 's. So, the following section will describe our methodology to obtain g_i 's and to adjust the diffusion model also.

5 Adjusted diffusion model with Gaussian density

In general, there is no clear way to find the exact base distribution of $X(t)$. However, we could characterize the asymptotic distribution for the multi-server queues from the literature. Many research studies on multi-server queues have shown that the limit processes of the multi-server queues are Gaussian processes, and the empirical density functions of them are also close to the Gaussian density. Listing some of those, for the time-homogeneous multi-server queues, Iglehart [8] and Whitt [22] show weak convergence to the Ornstein-Uhlenbeck (OU) process, and Halfin and Whitt [6] proves weak convergence to Brownian motion and the OU process depending on the traffic.

Therefore, for a given t , *weak convergence provides the Gaussian distribution which is asymptotically true*. For the time-varying multi-server queues (with abandonments and retrials), as depicted in Figure 5, Mandelbaum and Pats [11] and Mandelbaum et al. [13] show that the empirical density is close to the Gaussian density. Furthermore, *the result in Mandelbaum et al. [13] implies the limit process is a Gaussian process if the fluid model hits the critically loaded time points for a countable number of times, which is true for our model*. Therefore, for our model, it is reasonable to utilize the Gaussian distribution as a base distribution to identify g_i 's since the Gaussian assumption is asymptotically true. Once we decide to use the Gaussian density, it provides following two additional




Figure 5: Empirical density vs Gaussian density

benefits:

1. The Gaussian distribution can be completely characterized by the mean and covariance matrix which can be obtained from the fluid and diffusion models.
2. By using Gaussian density, g_i 's can achieve smoothness even if f_i 's are not smooth, which enables us to apply Theorem 2 without additional assumptions.

The second benefit is not obvious and hence we provide a proof of that.

Lemma 3. *Let g_i 's be the rate functions of $Z(t)$ obtained from the Gaussian density. Then, g_i 's are differentiable everywhere.*

Proof. Define

$$\phi(x, y) = \frac{1}{(2\pi)^{n/2} |\Sigma|^{1/2}} \exp \left(- \frac{(y - x)' \Sigma^{-1} (y - x)}{2} \right).$$

Using Gaussian density,

$$g_i(t, x) = \int_{\mathbf{R}^d} f_i(t, y) \phi(x, y) dy.$$

For $j \in \{1, \dots, d\}$, since $\phi(x, y)$ is differentiable with respect to x_j and $|f_i(t, y) \frac{d}{dx_j} \phi(x, y)|$ is integrable,

$$\begin{aligned} \frac{d}{dx_j} g_i(t, x) &= \frac{d}{dx_j} \int_{\mathbf{R}^d} f_i(t, y) \phi(x, y) dy \\ &= \int_{\mathbf{R}^d} f_i(t, y) \frac{d}{dx_j} \phi(x, y) dy \quad \text{by applying Theorem 2.27 in Folland [4],} \end{aligned} \quad (17)$$

where x_j is j^{th} component of x . Therefore, g_i is differentiable with respect to x_j . □

Now, we have $g_i(\cdot, \cdot)$'s which are differentiable. Then, we can apply Theorem 2 to obtain the diffusion model for $Z_n(t)$. Note that similar to the adjusted fluid model, once we have the distribution of $X(t)$, the adjusted diffusion model is applicable to more general cases. Therefore, we first follow the notation in Section 3.1 and will come back to our multi-server queues with abandonments and retrials.

Proposition 2 (Adjusted diffusion model). *Let $g_i(\cdot, \cdot)$'s be the rate functions in $Z(t)$ obtained from Gaussian density. Define a sequence of scaled centered processes $\{V_n(t)\}$ for $t \leq T$ to be*

$$V_n(t) = \sqrt{n} (Z_n(t) - \bar{Z}(t)),$$

where $Z_n(t)$ and $\bar{Z}(t)$ are solutions to equations (11) and (15) respectively. If $f_i(t, x)$'s and $F(t, x)$ satisfy equations (13) and (14) respectively, then $\lim_{n \rightarrow \infty} V_n(t) = V(t)$, where

$$V(t) = \sum_{i=1}^k l_i \int_0^t \sqrt{g_i(s, \bar{Z}(s))} dW_i(s) + \int_0^t \partial G(s, \bar{Z}(s)) ds,$$

$W_i(\cdot)$'s are independent standard Brownian motions, and $\partial G(t, \bar{Z}(t))$ is the gradient matrix of $G(t, \bar{Z}(t))$ with respect to $\bar{Z}(t)$. Furthermore, $V(t)$ is a Gaussian process.

Proof. From definition of $G(t, x)$ in (12), we can easily verify that $G(t, x)$ is differentiable by Lemma 3 and hence $|G(t, x) - G(t, y)| \leq M|x - y|$ implies

$$\left| \frac{\partial}{\partial x_i} G(t, x) \right| \leq M_i \quad \text{for some } M_i < \infty, t \leq T, \text{ and } i \in \{1, \dots, d\}.$$

Therefore, by Theorem 2, we prove this proposition. $\square$

Corollary 3. *If f_i 's are constants or linear combinations of the components of $X(t)$. Then,*

$$X(t) = Z(t) \quad \text{in distribution.}$$

Proof. Using the linearity of expectation, we can verify $g_i(t, x) = f_i(t, x)$ for $i \in \{1, \dots, k\}$. $\square$

Finally, we have the adjusted fluid and diffusion models by utilizing Gaussian density. Therefore, instead of assuming measure zero at a set of non-differentiable points (as done in Mandelbaum et al. [13]), we compare the adjusted models with the empirical mean and covariance matrix. Note when we explain Theorem 5, we do not consider $\Sigma(t)$, the covariance matrix of $X(t)$. However, from Gaussian density, we know that $\Sigma(t)$ characterizes the base distribution and it can be obtained from Proposition 2. Therefore, we rewrite g_i 's to be functions of t , $\bar{Z}(t)$, and $\Sigma(t)$; i.e.

$$g_i(t, \bar{Z}(t)) \rightarrow g_i(t, \bar{Z}(t), \Sigma(t)) \quad \text{for } i \in \{1, \dots, k\} \text{ and} \quad (18)$$

$$G(t, \bar{Z}(t)) \rightarrow G(t, \bar{Z}(t), \Sigma(t)). \quad (19)$$

Proposition 3 (Mean and covariance matrix). *Let $Y(t) = \bar{Z}(t) + V(t)$. Then,*

$$E(Y(t)) = \bar{Z}(t) \quad \text{and} \quad (20)$$

$$\text{Cov}(Y(t), Y(t)) = \text{Cov}(V(t), V(t)) = \Sigma(t). \quad (21)$$

The quantities $\bar{Z}(t)$ and $\Sigma(t)$ are obtained by solving the following simultaneous ordinary differential

equations with initial values given by $\bar{Z}(0) = x_0$ and $\Sigma(0) = 0$:

$$\frac{d}{dt}\bar{Z}(t) = \sum_{i=1}^k l_i g_i(t, \bar{Z}(t), \Sigma(t)), \quad (22)$$

$$\frac{d}{dt}\Sigma(t) = A(t)\Sigma(t) + \Sigma(t)A(t)' + B(t)B(t)', \quad (23)$$

where $A(t)$ is the gradient matrix of $G(t, \bar{Z}(t), \Sigma(t))$ with respect to $\bar{Z}(t)$, and $B(t)$ is the $d \times k$ matrix such that its i^{th} column is $l_i \sqrt{g_i(t, \bar{Z}(t), \Sigma(t))}$.

Proof. Since $V(0) = 0$, from Corollary 1, we have (20) and (21). By rewriting (15) in Theorem 5 as a differential equation form, we have (22), and by Theorem 3, we have (23). Note that since both $\bar{Z}(t)$ and $\Sigma(t)$ are variables, we should solve (22) and (23) simultaneously. $\square$

Eventually, we now have the adjusted fluid and diffusion models for the general cases, and it is the time to return to our system as given in Section 2. Using Gaussian density, we can obtain the new rate functions, g_i 's, which correspond to f_i 's as follows.

$$\begin{aligned} g_1(t, x) &= \lambda_t, \\ g_2(t, x) &= \mu_t^2 x_2, \\ g_3(t, x) &= \mu_t^1 (n_t + (x_1 - n_t)\Phi(n_t, x_1, \sigma_{1_t}) - \sigma_{1_t}^2 \phi(n_t, x_1, \sigma_{1_t})), \\ g_4(t, x) &= \beta_t(1 - p_t) \left((x_1 - n_t)(1 - \Phi(n_t, x_1, \sigma_{1_t})) + \sigma_{1_t}^2 \phi(n_t, x_1, \sigma_{1_t}) \right), \quad \text{and} \\ g_5(t, x) &= \beta_t p_t \left((x_1 - n_t)(1 - \Phi(n_t, x_1, \sigma_{1_t})) + \sigma_{1_t}^2 \phi(n_t, x_1, \sigma_{1_t}) \right), \end{aligned}$$

where $\Phi(a, b, c)$ and $\phi(a, b, c)$ are function values at point a of the Gaussian CDF and PDF respectively with mean b and standard deviation c .

Since $f_1(t, x)$ and $f_2(t, x)$ are constant and linear with respect to x respectively, $g_1(t, x) = f_1(t, x)$ and $g_2(t, x) = f_2(t, x)$. The derivation of other $g_i(\cdot, \cdot)$'s is straightforward but requires some computational efforts and hence we provide the details in Appendix B. Note g_3 , g_4 , and g_5 include σ_{1_t} which is currently treated as a function of t but is used by the adjusted diffusion model (see equations (18) and (19)). With the g_i 's above, by Proposition 3, we finally obtain $E[Z(t)]$ and $Cov[Z(t), Z(t)]$ for $t \leq T$ and will use them to approximate the mean and covariance matrix of our

original process $X(t)$ in equations (1) and (2).

Although we obtain the functions g_i 's for our adjusted models, we need some intuition regarding how g_i 's contribute to increasing accuracy especially in the critically loaded phases. Thus, in the next section, we revisit the inaccuracy in the previous approaches and explain how our adjusted models treat this.

6 Discussion on function g_i 's

In this section, we are going to investigate the functions g_i 's precisely. In order to get a clearer intuition, we consider a simple $M_t/M_t/n_t$ queue which is a special case of our original model ($\beta_t = 0$, and $\mu_t^1 = \mu_t$). Let $x(t)$ denote the number of customers in the system at time t . Then, $x(t)$ is the solution to the following integral equation:

$$x(t) = x(0) + Y_1\left(\int_0^t \lambda_s ds\right) - Y_2\left(\int_0^t (x(s) \wedge n_s) \mu_s ds\right).$$

Here, for convenience, define $f_1(t, x) = \lambda_t$, $f_2(t, x) = (x \wedge n_t) \mu_t$, and $F(t, x) = \lambda_t - (x \wedge n_t) \mu_t$. Applying theorems in Section 3.1, we have the fluid model $\bar{x}(t)$ and diffusion model $u(t)$ from the following integral equations:

$$\begin{aligned} \bar{x}(t) &= x(0) + \int_0^t \lambda_s - (\bar{x}(s) \wedge n_s) \mu_s ds, \text{ and} \\ u(t) &= u(0) + \int_0^t \left(\sqrt{\lambda_s}, \sqrt{(\bar{x}(s) \wedge n_s) \mu_s} \right) \begin{pmatrix} dW_1(t) \\ dW_2(t) \end{pmatrix} ds + \int_0^t \partial F(s, \bar{x}(s)) ds, \end{aligned}$$

where

$$\partial F(t, \bar{x}(t)) = \begin{cases} -\mu_t & \text{if } \bar{x}(t) \leq n_t, \\ 0 & \text{otherwise.} \end{cases}$$

Notice that the drift part $\partial F(t, \bar{x}(t))$ of the diffusion model is completely determined by the fluid model and here we might encounter a serious problem. Suppose we observe several realizations of this multi-server queue. When the $\bar{x}(t)$ is much smaller than the number of server n_t (underloaded phase), then there is not great possibility that an observed process is overloaded or critically loaded.

Therefore, the drift part $-\mu_t$ is valid in that sense. Now, assume that $\bar{x}(t)$ is smaller than but fairly close to n_t . Then, it is likely that significant fraction of the realizations could be overloaded or critically loaded. However, the drift part is still $-\mu_t$ since *the possibility of being overloaded or critically loaded is completely ignored by the fluid model*. Furthermore, imagine $\bar{x}(t)$ now becomes slightly larger than n_t . Then, the drift part suddenly changes to zero. As a result, if $\bar{x}(t)$ is fluctuating close to n_t , i.e. *lingering*, then the drift part of the diffusion model would repeat sudden changes between the values $-\mu_t$ and 0. Undoubtedly, it produces sharp spikes in the diffusion model as shown in Figure 3 and make the quality of the approximation worse especially near the critically loaded phase.

Now, we turn our attention to the functions g_i 's. In Section 4, g_i 's in the adjusted fluid model would improve the accuracy in estimating the mean values of the system states. Then, one may ask a question how g_i 's affect the estimation accuracy of the covariance matrix. To answer the question, let us follow the procedure to obtain $g_2(t, \cdot)$. Note $g_1(t, \cdot) = f_1(t, \cdot)$.

Define $G(t, x) = g_1(t, x) - g_2(t, x) = \lambda_t - g_2(t, x)$. For a fixed t_0 , let $x = x(t_0)$, $\mu = \mu_{t_0}$, $n = n_{t_0}$ and $z = E[x(t_0)]$. Then,

$$g_2(t_0, z) = E[\mu(x \wedge n)] = \mu \left\{ E[x \mathbb{I}_{x \leq n}] + n \Pr[x > n] \right\}. \quad (24)$$

From equation (24), we could notice the following characteristics of the function $g_2(\cdot, \cdot)$.

1. If $\Pr[x > n] \rightarrow 1$, $g_2(t_0, z) \rightarrow \mu n$.
2. If $\Pr[x > n] \rightarrow 0$, $g_2(t_0, z) \rightarrow \mu z$.


Note that $\partial G(t, \bar{z}(t))$ changes smoothly over time between $-\mu$ and 0 according to $\Pr[x(t) > n_t]$ as $\Pr[x(t) > n_t]$ changes smoothly under our Gaussian assumption (in fact, any distribution having a differentiable density works). Therefore, even if the adjusted fluid model $\bar{z}(t)$ is lingering in the vicinity of n_t , the drift part of the adjusted diffusion model changes smoothly over time. In the following section, we provide several experimental results and show the effectiveness of the adjusted models.

7 Numerical results


We compare our adjusted models against the fluid and diffusion models with the measure-zero assumption in Mandelbaum et al. [13] for multi-server queues with abandonments and retrials. Under the similar settings in Mandelbaum et al. [13], we use 5,000 independent simulation runs and compare the simulation result with both methodologies. We use the constant rates for the parameters except the arrival rate. The arrival rate alternates between 45 and 55 every two time units. Figures 6 and 7 show the estimation of mean values from one experiment. The number of servers (n_t) is 50 and the service rate of each server is 1. As seen in Figure 6, the number of customers in service node ($x_1(t)$) stays near the critically loaded point for a long time. As Mandelbaum et al. [13] points out, the fluid model with the measure-zero assumption shows significant estimation errors for $E[x_2(t)]$. On the other hand, our adjusted fluid model provides excellent approximation results. Especially, one can recognize remarkable improvement in the estimation of $E[x_2(t)]$. For the mean value of $x_1(t)$, our adjusted fluid model provides a lot better approximation result than the method with the measure-zero assumption.

When we see the covariance matrix, we also notice our adjusted diffusion model shows dramatic improvement against the diffusion model with the measure-zero assumption. As seen in Figure 7, the diffusion model assuming measure zero causes “spikes” as also pointed out in Section 3.2. Our proposed model, however, provides excellent accuracy without spikes at all.

Besides this specific example, in order to verify the effectiveness of our methodology, we conduct several experiments with different parameter combinations. Table 1 describes the setting of each experiment. In Table 1, “svrs” is the number of servers (n_t), “alter” is the time length for which each arrival rate lasts, and “time” is the end time of our analysis. We already recognize that the method assuming measure zero works well when it *does not linger* too long near the non-differentiable points. For comparison, therefore, our experiments contain several cases where the system *does linger* relatively long around those points as well as the cases where it does not. Experiments 1-4 are intended to see the effects of *lingering* around the critically loaded points. We change $\beta_t = \beta$ and $p_t = p$ as well as the arrival rates in experiments 5-8 to see the effects of other parameters. In fact, from the other experiments not listed in Table 1, it turns out that changing other parameters does not affect estimation accuracy significantly. Experiments 9 and 10 are set to observe how larger




(a) Mean numbers by assuming measure zero




(b) Mean numbers by our proposed method

Figure 6: Comparison of mean values, $E[X(t)]$



(a) Covariance matrix by assuming measure zero



(b) Covariance matrix by our proposed method


Figure 7: Comparison of covariance matrix entries, $Cov[X(t), X(t)]$

Table 1: Experiments setting


exp #	svrs	λ_1	λ_2	μ_1	μ_2	β	p	alter	time
1	50	40	80	1	0.2	2.0	0.5	2	20
2	50	40	60	1	0.2	2.0	0.5	2	20
3	100	80	120	1	0.2	2.0	0.7	2	20
4	100	90	110	1	0.2	2.0	0.7	2	20
5	50	40	80	1	0.2	1.5	0.7	2	20
6	50	40	60	1	0.2	1.5	0.7	2	20
7	50	45	55	1	0.2	2.0	0.5	2	20
8	100	95	105	1	0.2	2.0	0.5	2	20
9	150	140	160	1	0.2	2.0	0.5	2	20
10	150	100	190	1	0.2	2.0	0.5	2	20

arrival rates and number of servers affect the estimation accuracy along with the lingering effect by increasing both of them.

Here we explain the overall results: for the details of numerical results, see Table 2-6 in Appendix C. Similar to the results in Figures 6 and 7, we observe that lingering does debase the quality of approximations significantly when assuming measure zero. On the other hand, we see that our proposed models provide excellent accuracy for both mean and covariance matrix. Even if we increase both arrival rates and number of servers, we notice that lingering still affects the estimation accuracy significantly when assuming measure zero but it does not in our models. Figure 8 illustrates the average percentile difference of both methods against the simulation. Figure 8 (a) is obtained by averaging all differences in the tables (Appendix C) across time. From Figure 8 (a), we notice that our proposed method shows promise relative to the method assuming measure zero. However, in order to clearly see the effectiveness of our proposed methodology, we select the experiments 2, 4, 6, 7, 8, and 9 where lingering near the critically loaded phase occurs. We graph the differences at a critically loaded time point for those. Since the average differences are obtained from our limited experiments, it does not provide an absolute comparison between two methods. Nonetheless, we can notice that our method provides accurate estimation results consistently, but the method with measure-zero assumption results in vast inaccuracy. Note that, in Figure 8 (b), huge estimation difference, more than 300%, is observed when estimating $Cov[x_1(t), x_2(t)]$ using the method with measure-zero. However, the graph is cropped at the 70% level for the illustration purpose.



(a) Average difference for all experiments



(b) Average difference at a critically loaded point

Figure 8: Average difference against simulation

8 Conclusion

In this paper, we initially explain the strong approximations used in the analysis of multi-server queues with abandonments and retrials and show potential problems that one faces in obtaining accuracy and computational tractability especially near the critically loaded phase. The first problem stems from the fact that expectation of a function of a random vector X is not equal to the value of the function of the expectation of X . Therefore, unless they are equal or close, the fluid model may not provide an accurate estimation of mean values of the system state. The second problem is caused by non-differentiability of rate functions which prevents applying the diffusion model in Kurtz [9] and causes significant estimation errors if we ignore it. Therefore, addressing these problems is quite important in order to develop accurate approximations as well as to achieve computational feasibility. For that, we proposed a methodology to obtain the exact estimation of mean values of system states and an approach to achieve computational tractability.

The basic idea of our approach is to construct a new stochastic process which has the fluid limit exactly same as the mean value of the system state. We proved that if rate functions in the original model satisfy the conditions to apply the fluid model, rate functions in the constructed model also satisfy those conditions. Therefore, we can apply the adjusted fluid model if we can apply the existing fluid model. It turns out that there is, in general, no computational method to obtain the adjusted fluid model exactly. Fortunately, there are several previous research studies that show the distribution of limit processes and empirical distributions are close to the Gaussian in multi-server queueing systems and hence we utilize Gaussian density to approximate it. By using Gaussian density, we see that rate functions in the constructed model are smooth and we are able to apply the diffusion model in Kurtz [9] even if we could not apply it to the original process.

To validate our proposed method, we provide several numerical examples. In the examples, we observe that our proposed method shows great accuracy compared with the fluid and diffusion approximations with measure-zero assumption (which is the only other way in the literature, to the best of our knowledge, that provides computational tractability). Due to space restriction, we have not shown all examples where our method works well. We, however, observe that in some other types of queues other than multi-server queues considered here, e.g. peer-to-peer networks, multi-class queues, the empirical density is not close to the Gaussian density. For those types of queues,

one can investigate the properties of specific rate functions that affect the shape of empirical density and can devise a new methodology to find the functions $g_i(\cdot, \cdot)$'s from other density functions in the future.

Acknowledgments

The authors would like to thank Dr. William A. Massey and Dr. Martin I. Reiman for their inputs and valuable discussions. This research was partially supported by NSF grant CMMI-0946935.

Appendices

A Proof of Lemmas 1 and 2

Proof of Lemma 1. To prove this lemma, we need to show that $E[|X(t)|] \leq K(1 + |E[X(t)]|)$ for $K < \infty$ and $t \leq T$. We first show it in the one-dimensional case and then extend it to the d -dimensional case.

Let, for fixed $t_0 \leq T$, $X = X(t_0)$ having mean μ and variance σ^2 , and $f_i(X) = f_i(t_0, X(t_0))$. Then, by Cauchy-Schwarz inequality,

$$E[|X|] \leq \sqrt{E[X^2]} = \sqrt{\mu^2 + \sigma^2} \leq |\mu| + \sigma \leq D(1 + |\mu|) \quad \text{for } D = \max(1, \sigma). \quad (25)$$

Now, we have the one-dimensional case and can move to the d -dimensional case. Suppose X has a mean vector μ and a covariance matrix Σ such that $X = (x_1, \dots, x_d)'$, $\mu = (\mu_1, \dots, \mu_d)'$. Then,

$$\begin{aligned} E[|X|] &= E\left[\sqrt{\sum_{i=1}^d x_i^2}\right] \leq E\left[\sum_{i=1}^d |x_i|\right] = \sum_{i=1}^d E[|x_i|] \\ &\leq D\left(d + \sum_{i=1}^d |\mu_i|\right) \quad \text{by (25) for } D = \max(1, \sigma_1, \dots, \sigma_d) \\ &\leq D\left(d + d\sqrt{\sum_{i=1}^d \mu_i^2}\right) \quad \text{by Cauchy-Schwarz inequality} \\ &= Dd(1 + |\mu|). \end{aligned} \quad (26)$$

Now we have $E[|X|] \leq K(1 + |E[X]|)$ for the d -dimensional random vector X where $K = Dd$. Then,

$$\begin{aligned} \left| E[f_i(X)] \right| &\leq E[|f_i(X)|] \leq C_i + C_i E[|X|] \quad \text{from assumption} \\ &\leq C_i + C_i K(1 + |\mu|) \leq D_i(1 + |\mu|) \quad \text{for } D_i = C_i + C_i K \quad \text{by equation (26)} \end{aligned}$$

Note $g_i(t_0, \mu) = E[f_i(X)]$. Since $|\Sigma|$ is bounded on $t \leq T$, if we make $t_0 > 0$ arbitrary, we prove the lemma. $\square$

Proof of Lemma 2. For fixed $t_0 \leq T$, let $X = X(t_0)$ and $Y = Y(t_0)$ and suppose X and Y have a same base distribution H_0 (we use H instead of F to avoid confusion with F in (4)) where $E[X] = \mu_1$ and $E[Y] = \mu_2$. Then, the distribution H_1 of X and H_2 of Y satisfy

$$\begin{aligned} H_1(x) &= H_0(x - \mu_1), \quad \text{and} \\ H_2(y) &= H_0(y - \mu_2), \end{aligned}$$

respectively. Now, we have

$$\left| E[F(X)] - E[F(Y)] \right| = \left| \int_{\mathbf{R}^d} F(x) dH_1 - \int_{\mathbf{R}^d} F(y) dH_2 \right|.$$

By transforming variables,

$$\begin{aligned} \left| E[F(X)] - E[F(Y)] \right| &= \left| \int_{\mathbf{R}^d} F(x + \mu_1) dH_0 - \int_{\mathbf{R}^d} F(y + \mu_2) dH_0 \right| \\ &= \left| \int_{\mathbf{R}^d} (F(x + \mu_1) - F(x + \mu_2)) dH_0 \right| \quad \text{by linearity,} \\ &\leq \int_{\mathbf{R}^d} \left| (F(x + \mu_1) - F(x + \mu_2)) \right| dH_0 \\ &\leq M \int_{\mathbf{R}^d} |\mu_1 - \mu_2| dH_0 = M|\mu_1 - \mu_2| \quad \text{by assumption.} \end{aligned}$$

Note $G(t_0, \mu_1) = E[F(X)]$ and $G(t_0, \mu_2) = E[F(Y)]$. Then, by making $t_0 > 0$ arbitrary, we prove the second part, i.e. if $|F(t, x) - F(t, y)| \leq M|x - y|$ then $|G(t, x) - G(t, y)| \leq M|x - y|$. We can prove the first part, i.e. if $|f_i(t, x) - f_i(t, y)| \leq M|x - y|$, then $|g_i(t, x) - g_i(t, y)| \leq M|x - y|$, in a

similar fashion and hence we have the lemma. $\square$

B Derivation of $g_i(t, x)$'s

For fixed $t_0 > 0$, let $n = n_{t_0}$, $\mu_1 = \mu_{t_0}^1$, $\beta = \beta_{t_0}$, $p = p_{t_0}$, $x_1 = x_1(t_0) \sim N(z_1, \sigma_1^2)$, and $x_2 = x_2(t_0) \sim N(z_2, \sigma_2^2)$. For $z = (z_1, z_2)'$, we have

$$\begin{aligned}
g_3(t_0, z) &= E[\mu_1(x_1 \wedge n)] = \mu_1 \left\{ E[x_1 \mathbb{I}_{x_1 \leq n}] + nPr[x_1 > n] \right\} \\
&= \mu_1 \left[\int_{-\infty}^n \frac{x}{\sqrt{2\pi}\sigma_1} \exp\left(-\frac{(x-z_1)^2}{2\sigma_1^2}\right) dx + nPr[x_1 > n] \right] \\
&= \mu_1 \left[\frac{-\sigma_1}{\sqrt{2\pi}} \int_{-\infty}^n -\frac{x-z_1}{\sigma_1^2} \exp\left(-\frac{(x-z_1)^2}{2\sigma_1^2}\right) dx + z_1Pr[x_1 \leq n] + nPr[x_1 > n] \right] \\
&= \mu_1 \left[-\sigma_1^2 \frac{1}{\sqrt{2\pi}\sigma_1} \exp\left(-\frac{(n-z_1)^2}{2\sigma_1^2}\right) + (z_1 - n)Pr(x_1 \leq n) + n \right].
\end{aligned}$$

Therefore, by making $t_0 > 0$ arbitrary, we have $g_3(t, x)$.

Note $g_4(\cdot, \cdot)$ and $g_5(\cdot, \cdot)$ are same except a constant part with respect to x . Therefore, it is enough to derive $g_5(\cdot, \cdot)$. We can show that

$$\begin{aligned}
g_5(t_0, z) &= E[\beta p(x_1 - n)^+] = \beta p \{ E[x_1 \vee n] - n \} \\
&= \beta p \left\{ E[x_1 \mathbb{I}_{x_1 > n}] + nPr[x_1 \leq n] - n \right\} \\
&= \beta p \left[\int_n^\infty \frac{x}{\sqrt{2\pi}\sigma_1} \exp\left(-\frac{(x-z_1)^2}{2\sigma_1^2}\right) dx + nPr[x_1 \leq n] - n \right] \\
&= \beta p \left[\frac{-\sigma_1}{\sqrt{2\pi}} \int_n^\infty -\frac{x-z_1}{\sigma_1^2} \exp\left(-\frac{(x-z_1)^2}{2\sigma_1^2}\right) dx + z_1Pr[x_1 > n] + nPr[x_1 \leq n] - n \right] \\
&= \beta p \left[\sigma_1^2 \frac{1}{\sqrt{2\pi}\sigma_1} \exp\left(-\frac{(n-z_1)^2}{2\sigma_1^2}\right) + (z_1 - n)Pr(x_1 > n) \right].
\end{aligned}$$

Therefore, by making $t_0 > 0$ arbitrary, we have $g_5(t, x)$.

C Numerical results for Section 7

Table 2: Estimation of $E[x_1(t)]$ over time; difference from simulation

Experiments		Time (t)									
#	type	6	7	8	9	10	11	12	13	14	15
1	proposed	6.52	0.98	-3.39	-1.07	-3.05	-0.40	0.91	0.25	-0.69	-0.01
	meas. 0	4.42	0.82	-3.63	-1.94	-3.60	-0.23	0.75	-0.15	-2.59	0.11
2	proposed	2.69	0.44	-3.13	-0.82	-1.08	-0.32	0.48	0.15	-0.46	-0.05
	meas. 0	3.35	-0.42	-2.92	-1.64	-1.18	-1.01	0.85	-0.44	-0.36	-0.60
3	proposed	2.33	0.28	-3.11	-1.01	-1.36	-0.39	0.10	-0.02	-1.68	-0.15
	meas. 0	2.34	-0.42	-2.67	-1.55	-1.49	-1.00	0.52	-0.49	-0.54	-0.53
4	proposed	1.18	0.14	-1.54	-0.30	-0.01	0.12	0.22	0.22	-0.10	-0.02
	meas. 0	0.65	-0.96	-1.98	-1.32	-0.94	-0.95	0.04	-0.64	-0.61	-0.94
5	proposed	7.04	1.36	-3.67	-0.69	-1.38	-0.57	0.80	0.23	-2.82	-0.63
	meas. 0	5.55	1.04	-3.20	-0.93	-1.31	-0.53	0.46	0.06	-1.22	-0.18
6	proposed	3.61	0.76	-3.05	-1.13	-0.67	0.18	1.12	0.20	-0.95	-0.25
	meas. 0	2.53	-0.07	-3.01	-1.72	-1.46	-0.43	0.60	-0.47	-1.57	-0.80
7	proposed	1.93	0.65	-1.06	-0.25	-0.63	0.17	0.12	-0.21	-0.65	-0.20
	meas. 0	0.50	-0.86	-2.07	-1.51	-1.04	-0.73	-0.47	-1.07	-0.63	-0.76
8	proposed	0.72	0.07	-0.46	0.04	-0.04	-0.14	0.42	-0.07	-0.48	-0.01
	meas. 0	0.04	-0.98	-1.40	-0.91	-0.57	-0.85	-0.13	-0.69	-0.73	-0.46
9	proposed	0.81	0.25	-0.96	-0.25	-0.11	-0.09	0.38	-0.06	-0.24	-0.02
	meas. 0	0.53	-0.50	-1.31	-0.88	-0.34	-0.61	0.17	-0.51	-0.06	-0.32
10	proposed	6.44	1.18	-4.73	-1.73	-2.21	-0.45	0.30	-0.01	-1.10	-0.11
	meas. 0	6.46	0.77	-3.83	-1.62	-2.84	-0.83	0.84	0.00	-2.77	-0.60

Table 3: Estimation of $E[x_2(t)]$ over time; difference from simulation

Experiments		Time (t)									
#	type	6	7	8	9	10	11	12	13	14	15
1	proposed	-2.00	3.50	2.36	-0.53	0.57	-1.00	-0.99	-0.30	-0.44	-0.76
	meas. 0	11.68	12.60	7.38	5.88	11.64	8.18	5.24	6.29	10.47	7.82
2	proposed	-2.22	2.71	1.90	-2.44	-0.94	-1.82	-0.91	-0.10	-0.38	-0.76
	meas. 0	45.00	53.07	33.49	37.12	41.73	44.51	31.55	37.48	40.57	43.21
3	proposed	-2.49	1.88	1.00	-3.58	-2.09	-3.32	-3.08	-3.01	-3.15	-4.02
	meas. 0	28.64	37.65	19.44	21.88	24.73	28.38	16.37	21.45	22.77	26.96
4	proposed	0.24	2.66	1.35	-1.68	-0.91	-0.19	0.25	0.45	0.02	-0.53
	meas. 0	67.95	69.81	47.03	51.81	56.69	59.66	45.16	50.75	54.48	57.27
5	proposed	-1.01	4.41	3.16	-0.05	1.55	0.57	-0.58	-0.07	-0.09	-1.63
	meas. 0	9.61	12.28	7.50	5.73	11.28	9.42	5.36	6.00	9.51	8.02
6	proposed	-2.63	2.48	2.23	-2.39	-1.32	-0.84	-0.12	1.04	0.89	-0.04
	meas. 0	44.23	51.84	32.45	35.11	39.27	43.72	31.00	35.94	38.83	41.83
7	proposed	0.33	3.42	3.00	1.01	0.70	0.25	0.41	0.71	0.27	-0.17
	meas. 0	78.08	78.96	60.84	64.86	69.59	71.19	59.15	63.20	67.42	68.95
8	proposed	2.81	3.03	2.40	1.45	1.29	0.58	-0.08	0.41	0.12	-1.11
	meas. 0	92.68	90.60	73.97	77.06	80.98	81.48	70.82	74.24	77.96	78.55
9	proposed	-0.86	1.25	1.44	-0.77	-0.19	-0.18	0.08	0.84	0.42	0.41
	meas. 0	80.15	79.90	57.59	62.03	67.09	68.91	55.09	59.98	64.19	66.14
10	proposed	-2.67	6.62	3.79	-2.50	-0.49	-2.18	-1.99	-1.35	-1.38	-1.21
	meas. 0	8.53	23.91	10.73	8.77	10.78	13.05	5.67	8.96	9.13	11.69

Table 4: Estimation of $Var[x_1(t)]$ over time; difference from simulation

Experiments		Time (t)									
#	method	6	7	8	9	10	11	12	13	14	15
1	proposed	6.94	0.94	-1.92	-2.02	-3.66	-0.20	1.70	-1.02	0.49	2.89
	meas. 0	-11.03	2.93	-1.93	17.31	-24.44	1.42	1.66	14.89	-19.73	4.16
2	proposed	2.84	3.83	-6.05	-0.10	-0.50	4.24	1.62	2.67	-1.02	1.62
	meas. 0	-6.28	16.69	6.76	-14.45	-12.97	17.62	12.90	-11.61	-14.51	15.29
3	proposed	4.15	2.09	-0.60	2.15	-6.57	0.50	-3.10	1.15	2.76	3.74
	meas. 0	-0.56	13.38	7.30	-8.74	-12.97	11.92	4.53	-10.16	-2.13	14.22
4	proposed	-0.52	-4.36	-2.81	3.07	-0.03	2.96	0.79	1.27	3.30	0.35
	meas. 0	-16.38	11.18	14.13	-12.86	-17.81	17.93	16.94	-15.33	-14.05	15.32
5	proposed	6.83	-0.22	-2.49	0.09	-1.67	-3.27	1.71	-4.14	-0.55	1.98
	meas. 0	-2.30	1.03	-1.69	10.07	-10.59	-1.97	1.43	5.09	-7.69	3.42
6	proposed	5.22	0.62	-6.25	-0.81	-4.32	-1.95	4.41	1.97	-0.61	4.93
	meas. 0	-1.19	7.72	1.39	-7.42	-11.70	5.61	10.28	-4.33	-7.73	12.15
7	proposed	2.91	-2.29	-1.04	0.92	0.21	0.18	3.14	-1.10	4.36	2.28
	meas. 0	-17.83	14.52	18.27	-16.55	-22.37	17.07	20.88	-18.77	-18.14	19.01
8	proposed	-1.79	0.65	-0.43	0.83	3.35	-0.71	3.63	2.10	1.85	0.72
	meas. 0	-26.38	16.44	21.26	-18.37	-22.66	16.73	23.72	-17.42	-25.80	18.25
9	proposed	0.62	-0.86	-0.83	3.53	3.36	5.09	1.52	1.71	2.73	-1.37
	meas. 0	-17.84	13.72	17.09	-14.12	-17.40	19.78	18.07	-16.57	-19.03	14.36
10	proposed	4.48	-0.32	-9.84	1.26	-4.24	3.37	1.32	1.00	0.22	1.15
	meas. 0	4.12	7.27	-6.22	-2.69	-5.55	10.87	3.68	-3.27	-2.12	8.98

Table 5: Estimation of $Cov[x_1(t), x_2(t)]$ over time; difference from simulation

Experiments		Time (t)									
#	type	6	7	8	9	10	11	12	13	14	15
1	proposed	-3.03	-3.27	4.75	3.10	-1.72	-3.63	0.39	-4.00	-4.15	0.91
	meas. 0	25.05	-3.88	4.43	-6.75	15.78	-4.69	-0.30	-11.23	4.26	-2.03
2	proposed	-6.76	7.23	2.87	-4.73	11.50	-0.47	6.09	5.48	5.86	4.82
	meas. 0	29.60	-6.12	-6.56	3.81	36.90	-13.05	-1.41	12.63	31.69	-5.53
3	proposed	-6.74	-2.24	4.53	-9.51	-28.97	-3.44	-2.57	-6.43	5.52	-3.76
	meas. 0	25.67	-15.55	0.47	6.26	6.46	-15.26	-6.35	8.68	31.03	-13.44
4	proposed	-0.01	-13.57	-8.53	-12.42	-9.73	-0.00	-10.28	-16.74	-1.43	-11.64
	meas. 0	58.61	-29.39	-21.29	10.14	44.87	-14.34	-22.28	5.51	46.98	-26.05
5	proposed	-7.19	-0.18	0.13	2.88	7.13	-8.20	-0.42	1.07	4.17	-1.88
	meas. 0	19.73	-4.03	-1.03	-12.97	26.24	-11.11	-1.32	-12.31	20.40	-4.25
6	proposed	2.91	4.90	2.63	-7.64	-7.99	1.17	2.14	3.80	6.94	7.92
	meas. 0	37.93	-15.54	-15.96	-3.32	25.88	-18.86	-15.00	6.43	34.79	-10.85
7	proposed	-4.38	-1.88	0.97	-14.36	3.15	-1.95	-1.19	-1.08	-4.77	-0.46
	meas. 0	52.91	-16.88	-16.53	-3.19	43.26	-15.54	-16.02	6.73	34.99	-12.29
8	proposed	-20.99	-4.21	-6.33	-4.51	3.12	-3.43	-1.85	-6.78	-1.81	-0.79
	meas. 0	64.94	-8.85	-22.74	13.30	51.79	-11.89	-15.66	7.80	44.16	-8.72
9	proposed	-15.01	-6.15	-6.27	3.33	-0.25	2.45	-6.00	-7.57	-6.93	-6.74
	meas. 0	55.84	-12.34	-17.97	19.55	45.76	-4.17	-15.42	7.67	37.97	-12.70
10	proposed	-18.70	7.57	-3.70	-4.76	8.09	-6.43	-2.86	-0.03	2.95	-1.11
	meas. 0	-21.43	-2.63	-5.67	-0.67	8.99	-15.71	-4.40	3.66	4.18	-9.87

Table 6: Estimation of $Var[x_2(t)]$ over time; difference from simulation

Experiments		Time (t)									
#	type	6	7	8	9	10	11	12	13	14	15
1	proposed	-2.15	3.52	1.48	-0.72	-0.34	-0.45	0.78	1.59	1.31	0.83
	meas. 0	6.74	5.31	2.34	-8.01	5.46	1.00	1.84	-2.78	2.81	-1.20
2	proposed	1.29	9.81	8.50	3.31	7.72	6.70	6.05	5.84	5.42	5.91
	meas. 0	7.06	14.88	-6.56	-0.09	17.07	12.12	-3.90	5.44	16.94	13.19
3	proposed	-5.60	2.13	-0.71	-5.14	-1.93	-3.21	-2.71	-1.18	-0.61	-0.79
	meas. 0	-2.22	4.53	-10.37	-5.14	4.01	1.00	-8.89	0.66	6.45	5.96
4	proposed	5.71	8.34	2.63	-2.01	-0.83	2.01	-0.03	-0.27	0.51	2.66
	meas. 0	28.49	26.24	-13.87	-3.06	15.93	15.77	-12.78	0.50	17.18	17.62
5	proposed	-0.97	4.30	1.25	-0.07	3.22	3.50	-0.22	-0.16	1.50	0.76
	meas. 0	3.67	5.28	1.60	-4.35	7.92	6.18	1.65	-2.36	5.70	4.03
6	proposed	2.23	11.10	8.33	2.94	4.21	4.30	1.38	2.79	3.63	5.03
	meas. 0	11.13	21.61	-3.19	-0.25	12.83	14.35	-6.15	1.31	12.95	14.75
7	proposed	5.23	7.48	5.57	1.60	2.20	4.24	3.45	4.88	5.03	4.33
	meas. 0	33.03	27.56	-16.08	-4.39	21.67	19.11	-11.03	2.36	24.85	19.64
8	proposed	10.11	7.03	3.99	2.44	3.47	2.45	2.53	2.25	1.73	2.30
	meas. 0	62.03	46.52	-13.63	0.58	30.10	23.25	-13.94	-0.09	26.38	20.60
9	proposed	8.18	7.49	3.22	0.53	3.83	4.55	3.14	4.24	4.90	5.28
	meas. 0	39.93	31.88	-18.20	-4.36	22.39	18.66	-12.73	1.21	23.00	18.98
10	proposed	-0.34	12.31	5.05	-2.01	1.38	1.13	-3.01	-3.64	-4.35	-1.66
	meas. 0	-5.73	7.15	-1.91	-3.68	0.93	-2.52	-8.00	-4.34	-3.94	-5.72

References

- [1] Ludwig Arnold. *Stochastic Differential Equations: Theory and Applications*. Krieger Publishing Company, 1992.
- [2] Patrick Billingsley. *Convergence of Probability Measures*. A John Wiley & Sons, Inc., Publication, 1999.
- [3] Stewart N. Ethier and Thomas G. Kurtz. *Markov Processes: Characterization and Convergence*. A John Wiley & Sons, Inc., Publication, 1 edition, 1986.
- [4] Gerald B. Folland. *Real Analysis : Modern Techniques and Their Applications*. A John Wiley & Sons, Inc., Publication, 2 edition, 1999.
- [5] O. Garnet, Avi Mandelbaum, and Martin I. Reiman. Designing a Call Center with Impatient Customers. *Manufacturing & Service Operations Management*, 4(3):208–227, Jan 2002.
- [6] Shlomo Halfin and Ward Whitt. Heavy-Traffic Limits for Queues with Many Exponential Servers. *Operations Research*, 29(3):567–588, May 1981.

- [7] Robert C. Hampshire, Otis B. Jennings, and William A. Massey. A Time-Varying Call Center Design via Lagrangian Mechanics. *Probability in the Engineering and Informational Sciences*, 23(02):231–259, Jan 2009.
- [8] Donald L Iglehart. Limiting Diffusion Approximations for the Many Server Queue and the Repairman Problem. *Journal of Applied Probability*, 2(2):429–441, 1965.
- [9] Thomas G. Kurtz. Strong approximation theorems for density dependent Markov chains. *Stochastic Processes and their Applications*, 6(3):223–240, Feb 1978.
- [10] Avi Mandelbaum and Gennady Pats. State-dependent queues: approximations and applications. *Institute for Mathematics and Its Applications*, 71:239–282, 1995.
- [11] Avi Mandelbaum and Gennady Pats. State-Dependent Stochastic Networks. Part I: Approximations and Applications with Continuous Diffusion Limits. *The Annals of Applied Probability*, 8(2):569–646, May 1998.
- [12] Avi Mandelbaum, William A. Massey, and Martin I. Reiman. Strong approximations for Markovian service networks. *Queueing Systems*, 30(1-2):149–201, Jan 1998.
- [13] Avi Mandelbaum, William A. Massey, Martin I. Reiman, Alexander Stolyar, and Brian Rider. Queue Lengths and Waiting Times for Multiserver Queues with Abandonment and Retrials. *Telecommunication Systems*, 21(2-4):149–171, Jan 2002.
- [14] Avishai Mandelbaum and Sergey Zeltyn. Staffing Many-Server Queues with Impatient Customers: Constraint Satisfaction in Call Centers. *Operations Research*, 57(5):1189–1205, September-October 2009.
- [15] William A Massey and Ward Whitt. Uniform acceleration expansions for Markov chains with time-varying rates. *The Annals of Applied Probability*, 8(4):1130–1155, Nov 1998.
- [16] Guodong Pang and Ward Whitt. Heavy-traffic limits for many-server queues with service interruptions. *Queueing Systems*, 61(2-3):167–202, Mar 2009.
- [17] Anatolii A. Puhalskii and Martin I. Reiman. The multiclass GI/PH/N queue in the Halfin-Whitt regime. *Advances in Applied Probability*, 32(2):564–595, Jun 2000.

- [18] Ward Whitt. *Stochastic Process Limits*. Springer, 1 edition, 2002.
- [19] Ward Whitt. Efficiency-Driven Heavy-Traffic Approximations for Many-Server Queues with Abandonments. *Management Science*, 50(10):1449–1461, Oct 2006.
- [20] Ward Whitt. Fluid Models for Multiserver Queues with Abandonments. *Operations Research*, 54(1):37–54, January-February 2006.
- [21] Ward Whitt. Queues with Service Times and Interarrival Times Depending Linearly and Randomly Upon Waiting Times. *Queueing Systems*, 6(4):335–351, Dec 1990.
- [22] Ward Whitt. On the heavy-traffic limit theorem for GI/G/ ∞ queues. *Advances in Applied Probability*, 14(1):171–190, Mar 1982.
- [23] David Williams. *Probability with Martingales*. Cambridge University Press, 1991.
- [24] Sergey Zeltyn and Avi Mandelbaum. Call Centers with Impatient Customers: Many-Server Asymptotics of the M/M/n+ G Queue. *Queueing Systems*, 51(3-4):361–402, Jan 2005.